




## Urban Heat Stress Tracker



# Hyderabad



**Research direction:** Anumita Roychowdhury

**Authors:** Avikal Somvanshi and Sharanjeet Kaur



© 2024 Centre for Science and Environment

Material from this publication can be used, but with acknowledgement.

**Citation:** Avikal Somvanshi and Sharanjeet Kaur 2024, Urban Heat Stress Tracker: Hyderabad, Centre for Science and Environment, New Hyderabad.

**Published by**

**Centre for Science and Environment**

41, Tughlakabad Institutional Area

New Hyderabad 110 062

Phones: 91-11-40616000

Fax: 91-11-29955879

E-mail: [sales@cseinida.org](mailto:sales@cseinida.org)

Website: [www.cseindia.org](http://www.cseindia.org)



# Urban Heat Stress Tracker

# Hyderabad

# Overview

---

Heatwaves have become a staple of Indian summer due to the climate change. No region of the country is immune to this worsening phenomenon. States and cities are publishing their heat action plans to safeguard their populations from the dangerous heat exposures during heatwaves. These plans, while outlining the measures for emergency response and preparedness, also define the responsibilities of stakeholder departments in the event of a heatwave. These policy interventions assume significance at a time when heat and temperature trends are expected to worsen due to climate change and growing urbanization.

The relevance of these policy actions need to be understood against the rapidly changing global climate. The technical summary of the Intergovernmental Panel on Climate Change (IPCC), Working Group-I, Sixth Assessment Report (AR6 WG-I) notes that it is almost certain that the frequency and intensity of heat extremes and duration of heat waves have increased since 1950 and this will keep increasing even if global warming is stabilized at 1.5°C.<sup>1</sup> Combining climate change projections with urban growth scenarios, it can be said with very high confidence that future urbanization will amplify the projected increase in local air temperature.

With reference to urban centres, the IPCC Working Group-II, in its assessment (AR6 WG-II), also notes with confidence that hot extremes, including heat waves, have intensified in cities. It further notes that urban areas experience air temperatures that are several degrees warmer than surrounding areas, especially during the night. The urban heat island effect can add 2°C to local warming, reducing the adaptive capacity of cities and increasing the aforementioned risks.<sup>2</sup> This is due to reduced ventilation, heat trapping by closely-spaced tall buildings, heat generated directly from human activities, heat-absorbing properties of concrete and urban building materials, and limited vegetation. Infrastructure related to transportation, water, sanitation, energy and others has been compromised by extreme and slow-onset events, resulting in economic losses and disruption of services, impacting the well-being of people.

This emerging scientific evidence of the adverse impact of rising heat on urban populations builds the case for a city-specific heat management regime and the urgent implementation of heat action plans in cities. Such planning approaches also need to go much deeper than the immediate emergency response to help cope with specific heat events during summer and prevent heat lock-in. This is not only about summer action for public health protection but more sustained action throughout the year to heat proof the city and undertake heat mitigation, along with monitoring, to improve the overall adaptive thermal comfort of built structures and reduce energy and carbon intensity of built environment.

Such planning and intervention are possible if cities develop a tracking mechanism for annual and diurnal trends in temperature, humidity and the overall heat index to inform planning and implementation. Understanding the trend in heat and humidity patterns over time as well as during the day and night is necessary.

It is often noted that health emergency action considers the high daytime temperatures and not the nighttime temperatures and relative humidity. This overall trend poses risks to both public health and the energy security of the city, underlying the need to integrate this consideration into informing the heat action plan. The heat problem is not just about focusing on daily maximum temperatures crossing the 45°C benchmark—the standard focus during summer—but involves a much more complex set of indices.

Urban heat mitigation also requires more robust scientific tracking of key indicators—not just ambient heat and temperature, but also surface heat absorption and land surface temperatures, changing land-use, including vegetative cover and water bodies that are determinants to the heat island effect. This requires effective leveraging of the available satellite technology. Given advancements in technology, such data is available but needs policy integration.

It is equally important to track the various impacts of rising heat in the cities. The increasing heat is known to compromise the adaptive thermal comfort of people in cities and increase the demand for active cooling and use of mechanical cooling systems, including air conditioning which is an energy guzzler. This impacts the overall energy demand and energy security of the city and the region. Yet,



this dilapidating aspect of heat on a city's natural cooling abilities, including the rising trend in electricity demand to keep cool, is never tracked and considered for the active thermal management of cities.

This deeper conversation has to begin now because Hyderabad and several other states and cities have started developing their respective heat action plans.

In view of this, the Centre for Science and Environment has carried out this case study of select metro cities of India to analyse the trends in heat, humidity, land surface temperature and change in land use patterns to bring out the complex nature of heat management in cities. This detailed analysis of the heat scape of Hyderabad considers the time frame from 2001 to 2023.

This analysis has focused on the trends in day and night time temperature, humidity levels, seasonal variations, heat trends during day and night, trend in land surface temperature and trend in built-up area in the six megacities. Analysing these trends have provided deep insight into what is needed to inform the heat management practice in the city.

## Methodology and data

The study is based on comparative statistical analysis of temperature and the humidity condition observed in Hyderabad since 2001. The study's definition of summer is the period from March to August. It is further divided into pre-monsoon (March-May) and monsoon (June-August) as per IMD classification. This is based on publicly available datasets from various national and global agencies. Ambient temperature and humidity data have been sourced from Indian Meteorological Department (IMD) weather stations at Palam and Safdarjung. An average of the findings from these two weather stations is used to represent Hyderabad in this study. Heat Index computation has been done using the U.S. National Oceanic and Atmospheric Administration's (NOAA) formula. Complex geospatial calculations have been done in python and ArcGIS.

Moreover, freely accessible MODIS Land Science data from NASA Earth Observations has been used for seasonal and long term analysis of land surface temperature. For more granular analysis of heat and land use conditions on extremely hot days, satellite imagery data from the United States Geological Survey (USGS) Earth Explorer website has been used. Landsat 7 Enhanced Thematic Mapper Plus (ETM+) and Landsat 8 operational land imager/thermal infrared sensor (OLI/TIRS) satellite imagery were downloaded and used to analyse the land surface temperature, land use, land cover and Normalized Difference Vegetation Index (Green cover).

This city-level assessment focuses on changes in heat patterns over the years for the summer season, urban expansion over the years, and land surface temperature variation during the summer of 2003, 2013, and 2022. For Hyderabad, the later analysis is based on 10 May 2003, 29 May 2013, 14 May 2022, and 9 May 2023.

# Highlights

---

- 2024 March-April so far has been hotter (about 1°C) compared to average of 2014-23.
- Hyderabad's summertime has registered 0.9°C drop in decadal average ambient air temperature but the relative humidity has increased by 10 per cent between 2001-10 and 2014-23.
- High humidity is responsible for adding on average 1.5°C of heat stress to the city.
- Hyderabad has 30-90 days in summer when the daily ambient temperature exceeding 37°C mark.
- Pre-monsoon are thermally more uncomfortable than monsoon in Hyderabad. Average heat index during monsoon is about 3°C less than pre-monsoon.
- Hyderabad's is not cooling down at night at same rate as it used to do during 2001-10. But the diurnal cooling down of land surface temperature between daytime and nighttime is down by 13 per cent.
- Urban heat island phenomena is stronger at night than daytime in Hyderabad. During the daytime core of Hyderabad is 0.7°C cooler than its peripheries and peri-urban areas during the summer. But at night the core of Hyderabad is 1.9°C warmer than its peripheries and peri-urban areas
- There is direct co-relation between increase in built-up area and increase in urban heat stress. Built up area has increased from 20.6 per cent in 2003 to 44.0 per cent in 2023. Green cover has increased from 8.9 per cent in 2003 to 26.5 per cent in 2023. Increase in green cover shows impact on daytime temperatures but has no impact on nighttime temperature and increasing heat index in the city.




# Key findings

## Decadal trend in summertime heat

**Ambient air temperature in Hyderabad during summertime has cooled down over last two decades:** Recent few Hyderabad summers (March-August) have been less hotter than the average of first decade of the 21<sup>st</sup> century (2001-10). Decadal summertime average for Hyderabad in 2001-10 used to be 29.6°C on average which has decreased to 28.7°C in 2014-23; a drop of 0.9°C (see *Graph 1: Trend in summertime seasonal average ambient temperature in Hyderabad 2011-2023*).

**Graph 1: Trend in summertime seasonal average ambient temperature in Hyderabad 2011-2023**



**Note:** Summer is defined as the period from March to August. A city's weather profile is based on average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

**Nature of heat is changing in Hyderabad with significant increase in relative humidity during summer even though ambient air temperature has gone down:** Average Relative Humidity (RH) has significantly increased in the last 10 summers compared to 2001-10 average of 55.4 per cent. Hyderabad's last ten summers have been 10 per cent more humid on average compared to its 2001-10 average, in fact average RH of all summers from 2016 has been higher than 60 per cent (see *Graph 2: Trend in summertime seasonal relative humidity in Hyderabad 2011-2023*).


Hyderabad might have registered a significant increase in relative humidity level but it is located in one of the driest climatic zones of India. This significant jump in decadal relative humidity still doesn't



bring its overall humidity levels to coastal cities which are located in more humid climates. In fact this increased humidity in part lowers the impact of high ambient air temperatures in Hyderabad.

This combination of high heat and humidity can compromise the human body's main cooling mechanism: sweating. The evaporation of sweat from skin cools our bodies, but higher humidity levels limit this natural cooling. As a result, people can suffer heat stress and illness, and the consequences can even be fatal even at much lower ambient temperatures. Impact of this increasing humidity can be measured on human thermal comfort via means of Heat Index (HI). According to the U.S. National Weather Service, the heat index is a measure of how hot it really feels when humidity is factored in with the actual temperature. It is considered that a heat index of 41°C is dangerous to human health.

**Graph 2: Trend in summertime seasonal relative humidity in Hyderabad 2011-2023**




**Note:** Summer is defined as the period from March to August. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

**Heat Index has fallen like the ambient temperature in Hyderabad but impact of humidity is increasing on the city's thermal conditions:** Rise of relative humidity during summers of Hyderabad has had minimal impact on the city's heat index (HI). Hyderabad's summer HI average during 2001-10 used to be 30.7°C (impact of humidity: 1.1°C) which has decreased to 30.2°C (impact of humidity: 1.5°C) during 2014-23 (see *Graph 3: Trend in summertime seasonal average Heat Index in Hyderabad 2011-2023*). Overall, 0.5°C decrease in average HI but impact of humidity has increased by 0.4°C in the city's heat index.

**Graph 3: Trend in summertime seasonal average Heat Index in Hyderabad 2011-2023**






**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data uptill 30 August 2023.  
**Source:** CSE analysis of climatological data from IMD

**Hyderabad summer on average is marginally less hotter than it was in the first decade of the century:** Decadal RH average (2014-23) rose by 10.4 per cent in Hyderabad compared to 2001-10 average. This has marginally nullified the 3.1 per cent drop in the decadal ambient air temperature. As Hyderabad’s decadal average HI is down only by 1.9 per cent (see *Graph 4: Trend in decadal summertime heat in Hyderabad 2014-23 vs 2001-2010*).

**Graph 4: Trend in decadal summertime heat in Hyderabad 2014-23 vs 2001-2010**



**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on the average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data until 30 August 2023.




Source: CSE analysis of climatological data from IMD

**Number of days with high ambient temperatures are stable over the last two decades but days with dangerously high heat index are on a rise:** Hyderabad average used to have 57 days in a summer with high ambient temperature (37°C+) during 2001-10 but it registered just 41 days with such high temperatures in 2023 summer (see *Graph 5: Trend in days with 37°C+ daily maximum temperature in Hyderabad 2018-2023*). In fact, last four summers have registered lesser number of days with 37°C+ ambient temperature compared to the 2001-10 average.

But just looking at the daily maximum temperature figure is not a good measure of thermal discomfort and heat stress on the population as daily average temperature and humidity are critical to parameters as well. Human body is worse at handling humid heat than dry heat. If the heat index crosses the 41°C mark it is considered dangerous to human beings. In 2023 summer, Hyderabad registered zero days when the daily average heat index crossed the danger threshold of 41°C (see *Graph 6: Trend in days with 41°C+ daily heat index in Hyderabad 2018-2023*). Given the dry nature of the Hyderabad climate even during 2001-10 no days were registered where danger HI level was crossed. Only one such day has been registered during the study period of this analysis and it was in 2020 summer.

Additionally, it must be noted that in Hyderabad the majority of days with 37°C+ ambient temperature occur during pre-monsoon period (March-May) compared to monsoon period (June-August).


**Graph 5: Trend in days with 37°C+ daily maximum temperature in Hyderabad 2018-2023**



**Note:** Summer is defined as the period from March to August. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

Source: CSE analysis of climatological data from IMD

**Graph 6: Trend in days with 41°C+ daily heat index in Hyderabad 2018-2023**



**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on the average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

## Pre-monsoon vs monsoon heat


**Dry pre-monsoon period is getting muggier but impact of humidity is not significant in Hyderabad:** Summer can be divided into two distinct periods, i.e. pre-monsoon or dry heat period and monsoon or humid heat period. IMD defines pre-monsoon as March to May, while monsoon is considered from June to August. Naturally relative humidity is much lower during pre-monsoon compared to monsoon period. The study has found that average Relative Humidity (RH) has significantly increased for both pre-monsoon and monsoon period compared to 2001-10 average for Hyderabad. Last ten pre-monsoons have been on average 15 per cent more humid compared to 2001-10 average. Meanwhile monsoon humidity levels have risen by 8 per cent (see *Graph 7: Trend in relative humidity in Hyderabad 2011-2023 a. Pre-monsoon; b. Monsoon*).

Humidity has little impact on the pre-monsoon heat conditions of Hyderabad it has added on average (2014-23) 1.1°C in terms of heat index, which is higher than 2001-10 average of 0.6°C (see *Graph 8: Trend in impact of relative humidity on the ambient air temperature in Hyderabad 2011-2023 a. Pre-monsoon; b. Monsoon*).


During monsoon, humidity are much elevated compared to pre-monsoon which has a significant impact on the heat conditions. Any additional increase in RH levels can absolutely exacerbated the heat stress. This impact of humidity on Hyderabad’s heat index during 2001-10 was of 1.6°C which has increased to 1.7°C in 2014-23 (see *Graph 8: Impact of relative humidity on the ambient air temperature in Hyderabad 2011-2023 a. Pre-monsoon; b. Monsoon*).



**Graph 7: Trend in relative humidity in Hyderabad 2011-2023**  
**a. Pre-monsoon**




**b. Monsoon**




**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.  
**Source:** CSE analysis of climatological data from IMD

**Graph 8: Impact of relative humidity on the ambient air temperature in Hyderabad 2011-2023**  
**a. Pre-monsoon**



**b. Monsoon**




**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

**Both pre-monsoon and monsoon have cooled down in Hyderabad; though thermal distinction between monsoon and pre-monsoon is diminishing:** During 2001-10, the Heat Index used to fall between pre-monsoon and monsoon in Hyderabad by 3.1°C on average. This has decreased to 2.9°C during 2014-23 (see *Graph 9: Trend in number dangerously hot days in Hyderabad pre-monsoon vs monsoon 2011-2023*). Both pre-monsoon and monsoon on average got cooler by 0.7°C and 0.5°C compared to 2001-10 average respectively.

**Graph 9: Trend in number dangerously hot days in Hyderabad pre-monsoon vs monsoon 2011-2023**



**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

## Land surface heat and land use pattern


**Hyderabad is not cooling down at night:** During summers of 2001-10, the land surface temperature (LST) used to come down on average by 13.2°C from the daytime peak to nighttime low in Hyderabad. In the last ten summers (2014-23) the nighttime cooling has reduced to 11.5°C. This



translates to roughly 13 per cent reduction in diurnal cooling down (see *Graph 10: Trend in summertime diurnal land surface temperature changes in Hyderabad 2014-2023*). It must be noted that the nighttime cooling is getting even lesser than 11°C in the last few years.

Hot nights are as dangerous as midday peak temperatures. People get little chance to recover from daytime heat slaughter if temperatures remain high overnight, exerting prolonged stress on the body. A study published in the *Lancet Planetary Health* by a group of scientists from China, South Korea, Japan, Germany and the U.S. noted that the risk of death from excessively hot nights would increase nearly six-fold.<sup>1</sup> This prediction is much higher than the mortality risk from daily average warming suggested by climate change models.

**Graph 10: Trend in summertime diurnal land surface temperature changes in Hyderabad 2014-2023**



**Note:** Summer is defined as the period from March to August. \* Data uptill 30 August 2023.  
**Source:** CSE analysis of monthly MODIS Land Science data from NASA Earth Observations.


**Analysis of Hyderabad’s spatial heat-scape shows that its core is not cooling down at night at the same rate as its peri-urban region:** City cores are usually hotter than their surrounding peri-urban and rural areas as high population and built-up density traps and retains heat for longer duration. It is called the urban heat island phenomenon. Analysis of NASA satellite images shows that Hyderabad’s nighttime land surface temperature exhibit urban heat island formation. But Hyderabad’s daytime land surface temperature exhibits inverse of an urban heat island, i.e. the core of the city is cooler than its peri-urban during the daytime.

During the daytime core of Hyderabad is 0.7°C cooler than its peripheries and peri-urban areas during the summer. But at night the core of Hyderabad is 1.9°C warmer than its peripheries and peri-urban areas (see *Graph 11: Spatial variation in land surface temperature among the core city, outer city and peri-urban region of Hyderabad*). At night the peri-urban area cools down 12.3°C while the city core cools down only 9.7°C. So the city core is cooling down 2.6°C less than its peri-urban.

<sup>1</sup> Cheng He et al 2022. “The effects of night-time warming on mortality burden under future climate change scenarios: a modelling study”, *The Lancet Planetary Health*, Volume 6, Issue 8. [https://doi.org/10.1016/S2542-5196\(22\)00139-5](https://doi.org/10.1016/S2542-5196(22)00139-5)



**Graph 11: Spatial variation in LST among the core city, outer city and peri-urban region of Hyderabad**




**Note:** Based on average of 2018, 2019, 2020, 2021, 2022 and 2023 data. Summer is defined as March to August. \* Data uptill 30 August 2023.

**Source:** CSE analysis of monthly MODIS Land Science data from NASA Earth Observations.

Hyderabad have become more concertize in last two decades but green space has also increased significantly which seems to have blunted the rise in urban heat stress: Built up area has increased from 20.6 per cent in 2003 to 44.0 per cent in 2023. Green cover has increased from 8.9 per cent in 2003 to 26.5 per cent in 2023 (See *Graph 12: Change in land use pattern in Hyderabad in last two decades*).

**Graph 12: Change in land use pattern among megacities in last two decades**




**Note:** Summer heat wave months (May-June) are chosen to analyse the Normalized Difference Vegetation Index (NDVI) and urban expansion for each year.

**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.

**Land Use pattern change analysis:** Hyderabad saw an increase in its builtup area, with an expansion from 298.30 sq. km in 2003 to 636.58 sq. km in 2023, which depicts a substantial rise in the percentage share of the city’s geographical area from 20.6 per cent in 2003 to 44.0 per cent in 2023 (See *Map 1: Growth in Urban Built-up in Hyderabad during 2003, 2014 and 2023*). Significant and rapid urban expansion has happened in all directions, particularly in the Northwest zone.

**Impact of land surface changes on the distribution of land surface temperature:** In 2003, the average LST of Hyderabad was 39.5 °C. The outer peripheries of the city were particularly hot with temperatures surpassing 40°C. The dense settlement areas and open barren land in the Northwest and Southern regions of the city consistently recorded higher surface temperatures. The maximum LST was observed near the agricultural land in the North-west regions and stood at 47.05 °C. Water bodies and areas with dense green cover showed temperatures as low as 22.5°C even on an extreme heat day. In 2023, the average LST of Hyderabad was 30.2 °C, significantly cooler compared to 2003. On an extreme heat day (May 27, 2023), highest temperature were recorded at Rajiv Gandhi International Airport where LST reached 36.7 °C. Open/barren land near Drone Academy Nadirgul Airfield in the southern region, and the dense built-up area in the Northwest zone recorded surface temperatures between 35 °C to 36 °C. The lowest temperature recorded over water bodies and areas with dense green cover and it stood at 17.05 °C (See *Map 2: Variation in Land Surface Temperature over Hyderabad 2003, 2014 and 2022*).


**Map 1: Growth in Urban Built-up in Hyderabad during 2003, 2014, and 2023**



**Legend**  
 City Boundary     Built-up

**Note:** Urban expansion for each year – 2003, 2014, and 2023. The red color depicts the urban growth in the city.  
**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.

**Map 2: Variation in Land Surface Temperature over Hyderabad for 2003, 2014 and 2023**



*The maximum Land surface temperature reached 47.05 °C on May 28, 2003. The Southeast and Northwest part of the city records temperature above 42 °C.  
 On May 18, 2014, Jawahar Nagar in the Northeast region records LST above 36 °C and some of the area in the Northwest area observed with temperature above 38 °C. In 2023, the city maximum land surface temperature observed at 36.7 °C, the lowest of all the decades.*

**Land Surface Temperature (°C)**

	<27		30.1 - 33		36.1 - 38		40.1 - 42
	27.1 - 30		33.1 - 36		38.1 - 40		>42

**Note:** Summer heat wave months (May- June) are chosen to analyze the Land Surface Temperature (LST). The respective date of acquisition of the images are May 28, 2003, May 18, 2014, and May 27, 2023.

**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.