




## Urban Heat Stress Tracker



# Mumbai



**Research direction:** Anumita Roychowdhury

**Authors:** Avikal Somvanshi and Sharanjeet Kaur



© 2024 Centre for Science and Environment

Material from this publication can be used, but with acknowledgement.

**Citation:** Avikal Somvanshi and Sharanjeet Kaur 2024, Urban Heat Stress Tracker: Mumbai, Centre for Science and Environment, New Mumbai.

**Published by**  
**Centre for Science and Environment**

41, Tughlakabad Institutional Area

New Mumbai 110 062

Phones: 91-11-40616000

Fax: 91-11-29955879

E-mail: [sales@cseinida.org](mailto:sales@cseinida.org)

Website: [www.cseindia.org](http://www.cseindia.org)



## **Urban Heat Stress Tracker**

# **Mumbai**

# Overview

---

Heatwaves have become a staple of Indian summer due to the climate change. No region of the country is immune to this worsening phenomenon. States and cities are publishing their heat action plans to safeguard their populations from the dangerous heat exposures during heatwaves. These plans, while outlining the measures for emergency response and preparedness, also define the responsibilities of stakeholder departments in the event of a heatwave. These policy interventions assume significance at a time when heat and temperature trends are expected to worsen due to climate change and growing urbanization.

The relevance of these policy actions need to be understood against the rapidly changing global climate. The technical summary of the Intergovernmental Panel on Climate Change (IPCC), Working Group-I, Sixth Assessment Report (AR6 WG-I) notes that it is almost certain that the frequency and intensity of heat extremes and duration of heat waves have increased since 1950 and this will keep increasing even if global warming is stabilized at 1.5°C.<sup>1</sup> Combining climate change projections with urban growth scenarios, it can be said with very high confidence that future urbanization will amplify the projected increase in local air temperature.

With reference to urban centres, the IPCC Working Group-II, in its assessment (AR6 WG-II), also notes with confidence that hot extremes, including heat waves, have intensified in cities. It further notes that urban areas experience air temperatures that are several degrees warmer than surrounding areas, especially during the night. The urban heat island effect can add 2°C to local warming, reducing the adaptive capacity of cities and increasing the aforementioned risks.<sup>2</sup> This is due to reduced ventilation, heat trapping by closely-spaced tall buildings, heat generated directly from human activities, heat-absorbing properties of concrete and urban building materials, and limited vegetation. Infrastructure related to transportation, water, sanitation, energy and others has been compromised by extreme and slow-onset events, resulting in economic losses and disruption of services, impacting the well-being of people.

This emerging scientific evidence of the adverse impact of rising heat on urban populations builds the case for a city-specific heat management regime and the urgent implementation of heat action plans in cities. Such planning approaches also need to go much deeper than the immediate emergency response to help cope with specific heat events during summer and prevent heat lock-in. This is not only about summer action for public health protection but more sustained action throughout the year to heat proof the city and undertake heat mitigation, along with monitoring, to improve the overall adaptive thermal comfort of built structures and reduce energy and carbon intensity of built environment.

Such planning and intervention are possible if cities develop a tracking mechanism for annual and diurnal trends in temperature, humidity and the overall heat index to inform planning and implementation. Understanding the trend in heat and humidity patterns over time as well as during the day and night is necessary.

It is often noted that health emergency action considers the high daytime temperatures and not the nighttime temperatures and relative humidity. This overall trend poses risks to both public health and the energy security of the city, underlying the need to integrate this consideration into informing the heat action plan. The heat problem is not just about focusing on daily maximum temperatures crossing the 45°C benchmark—the standard focus during summer—but involves a much more complex set of indices.

Urban heat mitigation also requires more robust scientific tracking of key indicators—not just ambient heat and temperature, but also surface heat absorption and land surface temperatures, changing land-use, including vegetative cover and water bodies that are determinants to the heat island effect. This requires effective leveraging of the available satellite technology. Given advancements in technology, such data is available but needs policy integration.

It is equally important to track the various impacts of rising heat in the cities. The increasing heat is known to compromise the adaptive thermal comfort of people in cities and increase the demand for active cooling and use of mechanical cooling systems, including air conditioning which is an energy guzzler. This impacts the overall energy demand and energy security of the city and the region. Yet,



this dilapidating aspect of heat on a city's natural cooling abilities, including the rising trend in electricity demand to keep cool, is never tracked and considered for the active thermal management of cities.

This deeper conversation has to begin now because Mumbai and several other states and cities have started developing their respective heat action plans.

In view of this, the Centre for Science and Environment has carried out this case study of select metro cities of India to analyse the trends in heat, humidity, land surface temperature and change in land use patterns to bring out the complex nature of heat management in cities. This detailed analysis of the heat scape of Mumbai considers the time frame from 2001 to 2023.

This analysis has focused on the trends in day and night time temperature, humidity levels, seasonal variations, heat trends during day and night, trend in land surface temperature and trend in built-up area in the six megacities. Analysing these trends have provided deep insight into what is needed to inform the heat management practice in the city.

## Methodology and data

The study is based on comparative statistical analysis of temperature and the humidity condition observed in Mumbai since 2001. The study's definition of summer is the period from March to August. It is further divided into pre-monsoon (March-May) and monsoon (June-August) as per IMD classification. This is based on publicly available datasets from various national and global agencies. Ambient temperature and humidity data have been sourced from Indian Meteorological Department (IMD) weather stations at Palam and Safdarjung. An average of the findings from these two weather stations is used to represent Mumbai in this study. Heat Index computation has been done using the U.S. National Oceanic and Atmospheric Administration's (NOAA) formula. Complex geospatial calculations have been done in python and ArcGIS.

Moreover, freely accessible MODIS Land Science data from NASA Earth Observations has been used for seasonal and long term analysis of land surface temperature. For more granular analysis of heat and land use conditions on extremely hot days, satellite imagery data from the United States Geological Survey (USGS) Earth Explorer website has been used. Landsat 7 Enhanced Thematic Mapper Plus (ETM+) and Landsat 8 operational land imager/thermal infrared sensor (OLI/TIRS) satellite imagery were downloaded and used to analyse the land surface temperature, land use, land cover and Normalized Difference Vegetation Index (Green cover).

This city-level assessment focuses on changes in heat patterns over the years for the summer season, urban expansion over the years, and land surface temperature variation during the summer of 2003, 2013, and 2022. For Mumbai, the later analysis is based on 10 May 2003, 29 May 2013, 14 May 2022, and 9 May 2023.

# Highlights

---

- **2024 March-April so far has been similar thermally to the average of 2014-23.**
- **Mumbai's summertime has registered 0.6°C increase in decadal average ambient air temperature while the relative humidity has increased by 7 per cent between 2001-10 and 2014-23.**
- **High humidity is responsible for adding on average 5°C of heat stress to the city. Heat Index of the city has increased by 7 per cent.**
- **Both pre-monsoons and monsoons have gotten more thermally uncomfortable (by over 2°C) in Mumbai. Thermal distinction between monsoon and pre-monsoon has disappeared.**
- **Mumbai's is not cooling down at night at same rate as it used to do during 2001-10. But the diurnal cooling down of land surface temperature between daytime and nighttime is down by 24 per cent.**
- **Urban heat island phenomena is stronger at night than daytime in Mumbai. During the daytime core of Mumbai is 3.5°C cooler than its peripheries and peri-urban areas during the summer. But at night the core of Mumbai is 0.4°C warmer than its peripheries and peri-urban areas**
- **There is direct co-relation between increase in built-up area and increase in urban heat stress. Mumbai's built up area has increased from 38.4 per cent in 2003 to 52.1 per cent in 2023. Green cover has decreased from 35.8 per cent in 2003 to 30.2 per cent in 2023.**




# Key findings

## Decadal trend in summertime heat

**Ambient air temperature in Mumbai during summertime shows little change:** Recent few Mumbai summers (March-August) have been hotter than the average summer from the first decade of the 21<sup>st</sup> century (2001-10). Decadal summertime average for Mumbai is 29.2°C which is an increase of 0.6°C compared to 2001-10 (see *Graph 1: Trend in summertime seasonal average ambient temperature in Mumbai 2011-2023*).

**Graph 1: Trend in summertime seasonal average ambient temperature in Mumbai 2011-2023**



**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD


**Nature of heat is changing in Mumbai with significant increase in relative humidity during summer in addition to increase in ambient air temperature:** Average Relative Humidity (RH) in Mumbai has significantly increased in the last 10 summers compared to 2001-10 average. Mumbai’s last ten summers have been 7 per cent more humid on average compared to its 2001-10 average (see *Graph 2: Trend in summertime seasonal relative humidity in Mumbai 2011-2023*). This significant increase in relative humidity level compounds the thermal comfort issues of Mumbai which is located in humid climatic zone and already has a high RH baseline to start with.

This combination of high heat and humidity can compromise the human body’s main cooling mechanism: sweating. The evaporation of sweat from skin cools our bodies, but higher humidity levels limit this natural cooling. As a result, people can suffer heat stress and illness, and the consequences can even be fatal even at much lower ambient temperatures. Impact of this increasing humidity can be measured on human thermal comfort via means of Heat Index (HI). According to the U.S. National



Weather Service, the heat index is a measure of how hot it really feels when humidity is factored in with the actual temperature. It is considered that a heat index of 41°C is dangerous to human health.

**Graph 2: Trend in summertime seasonal relative humidity in Mumbai 2011-2023**




**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

**Heat Index rising faster than ambient temperature in Mumbai:** Given the rise of relative humidity during summer, the heat index (HI) has also risen. In the last 10 summers Mumbai’s HI average stood at 34.2°C (impact of humidity: 5.0°C) making the feel like temperature much hotter than ambient temperature. During 2001-10, the summer HI average was 31.9°C (see *Graph 3: Trend in summertime seasonal average Heat Index in Mumbai 2011-2023*). HI is a better measure of human thermal discomfort which accounts for the impact of both ambient heat and relative humidity.

**Graph 3: Trend in summertime seasonal average Heat Index in Mumbai 2011-2023**






**Note:** Summer is defined as the period from March to August. A city's weather profile is based on average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data uptill 30 August 2023.  
**Source:** CSE analysis of climatological data from IMD

**Mumbai summer on average is over 7 per cent hotter than it was in the first decade of the century:** Decadal RH average (2014-23) rose by 7.4 per cent in Mumbai compared to 2001-10 average. This is in addition to 1.9 per cent increase in the decadal ambient air temperature which has taken Mumbai's decadal average HI is up by 7.1 per cent (see *Graph 4: Trend in decadal summertime heat in Mumbai 2014-23 vs 2001-2010*).

**Graph 4: Trend in decadal summertime heat in Mumbai 2014-23 vs 2001-2010**



**Note:** Summer is defined as the period from March to August. A city's weather profile is based on the average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data until 30 August 2023.




**Source:** CSE analysis of climatological data from IMD

**Number of days with high ambient temperatures or dangerously high heat index are stable over the last two decades despite increase in overall heat stress:** Mumbai average used to have 5 days in a summer with high ambient temperature (37°C+) during 2001-10 but it registered just 4 days with such high temperatures in 2022 and 2023 summers (see *Graph 5: Trend in days with 37°C+ daily maximum temperature in Mumbai 2018-2023*).

But just looking at the daily maximum temperature figure is not a good measure of thermal discomfort and heat stress on the population as daily average temperature and humidity are critical to parameters as well. Human body is worse at handling humid heat than dry heat. If the heat index crosses the 41°C mark it is considered dangerous to human beings. During 2001-10, there were zero days when the danger HI mark of 41°C were breached in Mumbai but the summers of 2018 and 2019 registered 5 days and 2 days when the HI danger levels were crossed respectively (see *Graph 6: Trend in days with 41°C+ daily heat index in Mumbai 2018-2023*). Last four summers the 41°C HI danger level has not been breached.

Additionally, it must be noted that in Mumbai the days with 37°C+ ambient temperature occur exclusively during pre-monsoon period (March-May). Meanwhile, days with 41°C+ HI in Mumbai happen largely during monsoon period (June-August).


**Graph 5: Trend in days with 37°C+ daily maximum temperature in Mumbai 2018-2023**



**Note:** Summer is defined as the period from March to August. A city’s weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

**Graph 6: Trend in days with 41°C+ daily heat index in Mumbai 2018-2023**



**Note:** Summer is defined as the period from March to August. A city's weather profile is based on the average of all IMD weather stations located in the city. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

## Pre-monsoon vs monsoon heat


**Dry pre-monsoon period is getting muggier but the heat impact of humidity is most pronounced during monsoon:** Summer can be divided into two distinct periods, i.e. pre-monsoon or dry heat period and monsoon or humid heat period. IMD defines pre-monsoon as March to May, while monsoon is considered from June to August. Naturally relative humidity is much lower during pre-monsoon compared to monsoon period. This study has found that in Mumbai average Relative Humidity (RH) has significantly increased for both pre-monsoon and monsoon period compared to 2001-10 average. Last ten pre-monsoons have been on average 5 per cent more humid compared to 2001-10 average. Meanwhile monsoon humidity levels have risen by 9 per cent (see *Graph 7: Trend in relative humidity in Mumbai 2011-2023 a. Pre-monsoon; b. Monsoon*).

Humidity's impact on the pre-monsoon ambient heat conditions of Mumbai used to be 2.8°C during 2001-10 which has increased to 4.4°C during 2014-23, a staggering 56 per cent jump (see *Graph 8: Trend in impact of relative humidity on the ambient air temperature in Mumbai 2011-2023 a. Pre-monsoon; b. Monsoon*).


During monsoon, humidity are naturally elevated compared to pre-monsoon which has a significant impact on the heat conditions which translated to additional 3.8°C in terms of HI over ambient heat during 2001-10. The 9 per cent increase in the RH during 2014-23 monsoons translates to 5.7°C in terms of HI over ambient heat. An increase of 48 per cent in impact of RH on ambient heat (see *Graph 8: Impact of relative humidity on the ambient air temperature in Mumbai 2011-2023 a. Pre-monsoon; b. Monsoon*). This additional RH levels are exacerbating the heat stress in Mumbai.



**Graph 7: Trend in relative humidity in Mumbai 2011-2023**  
**a. Pre-monsoon**




**b. Monsoon**




**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.  
**Source:** CSE analysis of climatological data from IMD

**Graph 8: Impact of relative humidity on the ambient air temperature in Mumbai 2011-2023**  
**a. Pre-monsoon**




**b. Monsoon**



**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.  
**Source:** CSE analysis of climatological data from IMD

**Both pre-monsoons and monsoons have gotten more thermally discomfortable in Mumbai; distinction between monsoon and pre-monsoon is disappearing:** During 2001-10, the Heat Index used to rise between pre-monsoon and monsoon in Mumbai by 0.2°C on average. This has decreased to just 0.1°C during 2014-23. Even though the difference between pre-monsoon and monsoon has almost disappeared in terms of heat index but both parts of the summer are over 2°C compared to 2001-10(see *Graph 9: Decadal change in heat index in Mumbai pre-monsoon vs monsoon*).

**Graph 9: Decadal change in heat index in Mumbai pre-monsoon vs monsoon**



**Note:** Pre-monsoon refers to the months of March, April and June. Monsoon falls within June, July and August. Heat index has been calculated using the U.S. National Oceanic and Atmospheric Administration formula. A city's weather profile is based on the average of all IMD weather stations located in the city. \* Data until 30 August 2023.

**Source:** CSE analysis of climatological data from IMD

## Land surface heat and land use pattern


**Mumbai is not cooling down at night:** During summers of 2001-10, the land surface temperature (LST) used to come down on average by 10.2°C from the daytime peak to nighttime low in Mumbai. In



the last ten summers (2014-23) the nighttime cooling has reduced to 7.8°C. This translates to a staggering 24 per cent reduction in diurnal cooling down (see *Graph 10: Trend in summertime diurnal land surface temperature changes in Mumbai 2014-2023*). It must be noted that the nighttime cooling is getting even lesser in the last few years.

Hot nights are as dangerous as midday peak temperatures. People get little chance to recover from daytime heat slaughter if temperatures remain high overnight, exerting prolonged stress on the body. A study published in the *Lancet Planetary Health* by a group of scientists from China, South Korea, Japan, Germany and the U.S. noted that the risk of death from excessively hot nights would increase nearly six-fold.<sup>1</sup> This prediction is much higher than the mortality risk from daily average warming suggested by climate change models.

**Graph 10: Trend in summertime diurnal land surface temperature changes in Mumbai 2014-2023**



**Note:** Summer is defined as the period from March to August. \* Data uptill 30 August 2023.  
**Source:** CSE analysis of monthly MODIS Land Science data from NASA Earth Observations.



**Analysis of Mumbai’s spatial heat-scape shows that its core is not cooling down at night at the same rate as its peri-urban region:** City cores are usually hotter than their surrounding peri-urban and rural areas as high population and built-up density traps and retains heat for longer duration. It is called the urban heat island phenomenon. Analysis of NASA satellite images shows that Mumbai’s nighttime land surface temperature exhibit urban heat island formation. But Mumbai’s daytime land surface temperature exhibits inverse of an urban heat island, i.e. the core of the city is cooler than its peri-urban during the daytime.

During the daytime core of Mumbai is 3.5°C cooler than its peripheries and peri-urban areas during the summer. But at night the core of Mumbai is 0.4°C warmer than its peripheries and peri-urban areas (see *Graph 11: Spatial variation in land surface temperature among the core city, outer city and peri-urban region of Mumbai*). At night the peri-urban area cools down 10.6°C while the city core cools down only 6.7°C. So the city core is cooling down 3.9°C less than its peri-urban.

<sup>1</sup> Cheng He et al 2022. “The effects of night-time warming on mortality burden under future climate change scenarios: a modelling study”, *The Lancet Planetary Health*, Volume 6, Issue 8. [https://doi.org/10.1016/S2542-5196\(22\)00139-5](https://doi.org/10.1016/S2542-5196(22)00139-5)



**Graph 11: Spatial variation in LST among the core city, outer city and peri-urban region of Mumbai**



**Note:** Based on average of 2018, 2019, 2020, 2021, 2022 and 2023 data. Summer is defined as March to August. \* Data until 30 August 2023.


**Source:** CSE analysis of monthly MODIS Land Science data from NASA Earth Observations.

**Note:** Based on average of 2018, 2019, 2020, 2021, 2022 and 2023 data. Summer is defined as March to August. \* Data uptill 30 August 2023.

**Source:** CSE analysis of monthly MODIS Land Science data from NASA Earth Observations.

**Mumbai has become more concertize in last two decades which has contributed to rise in urban heat stress:** Mumbai's built up area has increased from 38.4 per cent in 2003 to 52.1 per cent in 2023. Green cover has decreased from 35.8 per cent in 2003 to 30.2 per cent in 2023 (See Graph 12: Change in land use pattern in Mumbai in last two decades).

**Graph 12: Change in land use pattern among megacities in last two decades**




**Note:** Summer heat wave months (May-June) are chosen to analyse the Normalized Difference Vegetation Index (NDVI) and urban expansion for each year.

**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.

**Land Use pattern change analysis:** Mumbai saw a significant increase in its built-up area, with an expansion from 241.15 sq. km in 2003 to 326.81 sq. km in 2023, which depicts a gradual rise in the percentage share of the city's geographical area from 38.4 per cent in 2003 to 52.1 per cent in 2023 (See *Map 1: Growth in Urban Built-up in Mumbai during 2003, 2013 and 2023*). Significant and rapid urban expansion has happened all around the city, especially in the East, South, Southwest, and South East zone.


**Impact of land surface changes on the distribution of land surface temperature:** In 2003, the average LST of Mumbai was 33.98 °C. Temperatures in excess of 38°C were recorded over highly dense built-up areas, barren land, and cropland mostly located the North, North-east, and South-east regions of the city. The highest land surface temperature was observed at Deonar dumping ground and it stood at 47.5 °C on an extreme heat day (April 22, 2003). Water bodies and areas with dense green cover showed temperatures as low as 25°C even on an extreme heat day. In 2023, the average LST of Mumbai was 32.07 °C, about 2°C cooler than 2003. However, the Deonar dumping ground in the Southeast region and cropland in the upper Northeast area, as well as the suburbs surrounding the Chhatrapati Shivaji International Airport in the central part of the city have continued to registering temperatures above 38°C. On May 31, 2023, the highest surface temperature was recorded over bare and open lands in the northeast zone of the city and it stood at 40.6°C. The lowest temperature recorded over water bodies and areas with dense green cover and it stood at 20.8 °C (See *Map 2: Variation in Land Surface Temperature over Mumbai for 2003, 2013 and 2022*).

**Map 1: Growth in Urban Built-up in Mumbai during 2003, 2013, and 2023**



**Note:** Urban expansion for each year – 2003, 2013, and 2023. The red color depicts the urban growth in the city.  
**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.

**Map 2: Variation in Land Surface Temperature over Mumbai for 2003, 2013 and 2023**



*On April 22, 2003, LST over the Doonar region in South-east part of the city and the agricultural area in the upper North-east region records temperature above 42 °C.  
 In 2013, the severe pockets of LST spread over the barren land near Radha Soami Satsang Beas in the northwest region, cropland in the North-east part and Deonar dumping ground in the south east recorded temperature above 40 °C.  
 On May 31, 2023, the hotspots remain consistent along with some isolated pockets in the city with temperature above 40 °C.*



**Note:** Summer heat wave months (April-May) are chosen to analyze the Land Surface Temperature (LST). The respective date of acquisition of the images are April 22, 2003, May 11, 2013, and May 31, 2023.  
**Source:** CSE analysis of Landsat 7 and Landsat 8 satellite images from United States Geological Survey (USGS) Earth Explorer.