




Air Quality Tracker **Ground-level Ozone**



Pune

Metropolitan Region



Research direction: Anumita Roychowdhury

Authors: Avikal Somvanshi and Sharanjeet Kaur



© 2024 Centre for Science and Environment

Material from this publication can be used, but with acknowledgement.

Citation: Avikal Somvanshi and Sharanjeet Kaur 2024, Air Quality Tracker
Ground-level Ozone: Pune Metropolitan Region, Centre for Science and Environment, New
Delhi.

Published by

Centre for Science and Environment

41, Tughlakabad Institutional Area

New Delhi 110 062

Phones: 91-11-40616000

Fax: 91-11-29955879

E-mail: sales@cseinida.org

Website: www.cseindia.org



Air Quality Tracker
Ground-level Ozone

Pune
Metropolitan Region

Overview

Centre for Science and Environment (CSE) has alerted from time to time about the growing problem of ground-level ozone in Indian cities. While policy and public attention is nearly fully drawn towards very high level of particulate pollution, the challenge of this highly toxic gas has not attracted adequate policy attention for mitigation and prevention. Inadequate monitoring, limited data and inappropriate methods of trend analysis have weakened the understanding of this growing public health hazard. This requires early action.

The summer of 2024 has witnessed widespread ground-level ozone exceedance making the air of Pune and its larger metropolitan region even more toxic. This summer the geographical spread of the problem is even more wide spread as compared to last summer. The toxic built up is also lasting longer at locations where it is happening this summer compared to previous summers.

Health evidence is also growing stronger. The 2020 State of Global Air report states that age-standardized rates of death attributable to ground-level ozone is among the highest in India and the seasonal 8-hour daily maximum concentrations have recorded one of the highest increases in India between 2010 and 2017– about 17 per cent. This requires deeper understanding of what is going on in different cities and regions to inform mitigation.

Due to the very toxic nature of ground-level ozone, the national ambient air quality standard for ozone has been set for only short-term exposures (one-hour and eight-hour averages), and compliance is measured by the number of days that exceed the standards. Compliance requires that the standards are met for 98 per cent of the time of the year. It may exceed the limits on two per cent of the days in a year, but not on two consecutive days of monitoring. In other words, there should not be more than eight days in a year when the ozone standard is breached, and none of those allowed exceedances can be on two consecutive days.

The standard practice of Central Pollution Control Board to average out the data of all stations in the city to determine daily AQI does not work for ground-level ozone as it is a short-lived and hyper-localised pollutant. A citywide average concentration level over an extended time frame does not indicate the severity of the problem and health implication from local build up and exposure for people living in hotspots.

Global experience shows that there is usually a trade-off. As particulate pollution is reduced the problem of NO_x and ground-level ozone increase. Globally, regulators are tightening the regulatory benchmark for ozone to address the toxic threat which – given its complex chemistry, is difficult to address. India should prevent this trap.

Why ozone needs special attention? Complex chemistry of ground-level ozone makes it a difficult pollutant to track and mitigate. Ground-level ozone is not directly emitted from any source. It is produced from complex interaction between nitrogen oxides (NO_x) and volatile organic compounds (VOCs) that are emitted from vehicles, power plants, factories, and other combustion sources and undergo cyclic reactions in the presence of sunlight to generate ground-level ozone. VOCs can also be emitted from natural sources, such as plants. Ozone not only builds up in cities but also drifts long distances to form a regional pollutant that makes both local and regional action necessary.

This highly reactive gas has serious health consequences. Those with respiratory conditions, asthma, chronic obstructive pulmonary disease, and particularly children with premature lungs and older adults are at serious risk. This can inflame and damage airways, make lungs susceptible to infection, aggravate asthma, emphysema, and chronic bronchitis and increase the frequency of asthma attacks leading to increased hospitalisation.

The investigation: This assessment has traced trends in ground-level ozone data from 2020 to 2024. The analysis is based on publicly available granular real time data (15-minute averages) from the CPCB's official online portal Central Control Room for Air Quality Management. The data has been captured from 48 official stations under the Continuous Ambient Air Quality Monitoring System (CAAQMS) spread across Pune Metropolitan Region. Pune (12) and Pimpri Chinchwad (3).

Given the volatile and highly localized nature of ground-level ozone pollution build-up and its variability across space, and consistent with the global good practice, this analysis has considered station level trends in terms of number of days exceeding the 8-hour standard over time. As ozone formation depends on complex atmospheric chemistry and on photochemical reaction its level varies across time and space horizon. Meteorological parameters such as sunny and warm weather, stagnant wind patterns etc have bearing on its formation. This analysis tracks exceedances at each station in core NCR. Breach of the standard by even one station is considered exceedance by the core NCR. Days with multiple stations exceeding the standard indicates the severity of the spatial spread and number of people exposed. Given that the data is capped at 200 µg/m³ by CPCB, it is not possible to determine how high the concentration really goes.


The study has considered global good practice and taken on board the USEPA approach of computing eight-hour averages for a day and then checking for the maximum value among them to capture the daily ozone pollution level. USEPA assesses city-wide or regional AQI based on the highest value recorded among all stations of the city or the region. Thus, trends have been calculated in terms of number of days when the daily level has exceeded the 8-hr standard (referred as exceedance days hereafter).

While analysing the data it has also been noted that the ozone data available on CPCB portal never exceeds 200 µg/m³, while data for the corresponding time on Delhi Pollution Control Committee may show higher levels. Therefore, due to this capping of data it is not possible to understand the nature of peaking in the city. This needs to be addressed as there are two sets of standard for ozone – 8-hourly standard of 100 µg/m³ and one hourly standard at 180 µg/m³. Capping makes assessment of one-hourly standard challenging.

Key findings

Ground-level ozone exceedance is reported on nearly half of this summer: This summer ground-level ozone exceedances were reported on 58 days between 1 April and 18 July. The worst days of ground ozone pollution were in the mid of May when 6 stations out of 16 stations of Pune Metropolitan Region reported exceedance (See *Graph 1: Daily variation in ground-level ozone exceedance in Pune Metropolitan Region*). The intensity of pollution was highest on 15 May when the regional average stood at $123 \mu\text{g}/\text{m}^3$.


Graph 1: Daily variation in ground-level ozone in Pune Metropolitan Region



Note: Based on exceedances recorded at the monitoring stations at Pune Metropolitan Region. Exceedance is computed as daily maximum 8-hr average crossing the ground-level ozone 8-hr standard, i.e. $100 \mu\text{g}/\text{m}^3$. Period of study is 1 April to 18 July 2024. Source: CSE analysis of CPCB realtime data.

Geographical spread of ground-level ozone pollution in Pune Metropolitan Region is on rise: Ground-level ozone usually exceeds the safety standard on all days of summer in some location in Pune Metropolitan Region every year. This year the spatial spread (number of stations exceeding the standard across Pune Metropolitan Region) has been 3.2 stations per day for period 1 January to 30 June. This is the highest in last five years. On an average only 1 station used to exceed the standard daily during the previous summers (See *Graph 2: Variation in average duration and spatial spread of daily exceedance in Pune Metropolitan Region*). One of the probable reason for the increase is the expansion of monitoring in the region in last couple of years.

Average duration of exceedance has also increased in last five years. This year so far, at the stations which reported exceedance it lasted on average 15.1 hours, it used to be below 13 hours in the previous years.

Graph 2: Variation in average duration and spatial spread of daily exceedance in Pune

Note: Based on exceedances recorded at the monitoring stations in Pune Metropolitan Region. Exceedance is computed as daily maximum 8-hr average crossing the ground-level ozone 8-hr standard, i.e. $100 \mu\text{g}/\text{m}^3$. Duration of exceedance is computed as number of hours the rolling 8-hr avg was exceeded at a station on a day. Period of the study is 1 Jan to 30 Jun. Source: CSE analysis of CPCB realtime data.


Savitribai Phule University and Pimpri Chinchwad are worst affected by ground-level ozone pollution: Savitribai Phule University in Pune is the most chronically affected by ground ozone pollution in Pune Metropolitan Region. It has exceeded the standard in this location for 119 days this year so far. It is followed by Gavalinagar and Thergaon in Pimpri as the worst polluted with 119 days and 99 days of exceedance respectively (See *Map 1: Hotspots of ground-level ozone exceedance in Pune Metropolitan Region* & *Table 1: Locations with most ground-level ozone exceedance in Pune Metropolitan Region*). Shivajinagar and Nigdi in Pune have least instances of ground-level ozone exceedances in the region.

Table 1: Locations with most ground-level ozone exceedance in Pune Metropolitan Region

SNb.	Station	Number of exceedance days
1	Savitribai Phule University, Pune	119
2	Gavalinagar, Pimpri	119
3	Thergaon, Pimpri	99
4	Park Street Wakad, Pimpri	33
5	Karve Road, Pune	22
6	Alandi, Pune	20
7	Mhada Colony, Pune	16
8	Bhosari, Pune	8
9	Shivajinagar, Pune	0
10	Nigdi, Pune	0
11	Hadapsar, Pune	0
12	Bhumkar Nagar, Pune	0
13	Katraj Dairy, Pune	0
14	Panchawati, Pune	0
15	MT, Pune	0

Note: Exceedance is computed as daily maximum 8-hr average crossing the ground-level ozone 8-hr standard, i.e. $100 \mu\text{g}/\text{m}^3$. Period of study is 1 January to 18 July 2024. Source: CSE analysis of CPCB realtime data.




Map 1: Hotspots of ground-level ozone exceedance in Pune Metropolitan Region



Note: Based on exceedances recorded at the monitoring stations in Pune Metropolitan Region. Exceedance is computed as daily maximum 8-hr average crossing the ground-level ozone 8-hr standard, i.e. $100 \mu\text{g}/\text{m}^3$. Period of study is 1 April to 18 July 2024.
Source: CSE analysis of CPCB realtime data.

Ground-level ozone hotspots are located in the areas with low levels of NO₂ and PM_{2.5}: The spatial distribution of ground-level ozone is inverse of the NO₂ and PM_{2.5} (see *Map 2: Spatial relationship among hotspots for key pollutants in Pune Metropolitan Region*). This bears out the fact that while ozone is created in polluted areas with nitrogen oxide being the catalyst, it also gets mopped up in high NO₂ areas as it further reacts. But the ozone that escapes to cleaner areas with less NO₂ builds up faster as unavailability of NO₂ hampers its dissipation. Park Street Wakad in Pimpri is exception to this phenomena as these stations report both high NO₂ and ground-level ozone.

Map 2: Spatial relationship among hotspots for key pollutants in Pune Metropolitan Region



Note: Seasonal average computed as mean of monthly averages based on daily 24-hr average for PM2.5 and NO₂, while daily maximum 8-hr average is used for ground-level ozone. Period of study is 1 April to 18 July 2024. Source: CSE analysis of CPCB realtime data.

Regional hourly ozone peak level is up by 200 per cent compared to lockdown times: Since CPCB caps the data at 200 $\mu\text{g}/\text{m}^3$ it is not possible to access precisely how high the ground-level ozone concentration can go up to, but for to get a relative understanding in this study hourly data averaged across all station and all days of April has been analysed. This indicative analysis shows that compared to April of 2020 ground-level ozone is not lingering in the air post sunset. Nevertheless, the hourly peak on an average has gone up by 200 per cent (See *Graph 3: Hourly cycle of ground level ozone and NO₂ in Pune Metropolitan Region – April 2020 v/s April 2024*). The re-emergence of morning and evening rush-hour traffic is helping




in neutralising ground-level ozone at sunrise and sunset as increased NO₂ levels cannibalise it.

The maximum 8-hour average was recorded at Savitribai Phule University in Pune when level hit 212.7 µg/m³ on 22 March 2024. It was followed by Gavalinagar in Pimpri (See *Table 2: Locations with the highest daily peak ground-level ozone pollution in Pune Metropolitan Region*).

Given the data cap of 200 µg/m³ enforced by CPCB at the 15-minute granularity, it is stunning to observe that multiple stations are still logging 8-hourly averages exceeding 180 µg/m³ every year. This underscores the magnitude of the pollution.

Graph 3: Hourly cycle of ground-level ozone and NO₂ in Pune Metropolitan Region – April 2020 v/s April 2024



Note: 24-hr profile is based on mean hourly concentration of ground-level ozone and NO₂ recorded at the monitoring stations in Pune Metropolitan Region for month of April in 2020 and 2023. Period of study is 1 April to 18 July 2024. Source: CSE analysis of CPCB realtime data.

Table 2: Locations with highest daily peak ground-level ozone pollution in Pune


SNo	Station	Highest daily 8-hr average in $\mu\text{g}/\text{m}^3$
1	Savitribai Phule University, Pune	212.7
2	Gavalinagar, Pimpri	185.2
3	Bhosari, Pune	163.6
4	Thergaon, Pimpri	163.5
5	Alandi, Pune	162.6
6	Mhada Colony, Pune	160.3
7	Park Street Wakad, Pimpri	141.0
8	Karve Road, Pune	138.2
9	Hadapsar, Pune	92.1
10	Shivajinagar, Pune	78.8
11	Bhunkar Nagar, Pune	77.5
12	Ngdi, Pune	74.2
13	Katraj Dairy, Pune	71.4
14	Panchawati, Pune	41.8
15	MT, Pune	12.7

Note: Based on daily maximum 8-hr average. Period of study is 1 January to 18 July 2024. Source: CSE analysis of CPCB realtime data.

Night-time ground-level ozone continues to persist: Ground-level ozone should ideally become negligible in the night air but MMR has been witnessing a rare phenomenon where ozone levels remain elevated hours after sunset. This was found to be very wide-spread in Delhi during the lockdowns of 2020 summers but it is now been noted in MMR this year as well. This night-time ozone was noted at 1.4 stations on average every night (See *Graph 4: Daily variation in occurrence of high night-time ground-level ozone pollution*). Five stations recorded nighttime ozone on multiple days this year during April and May. It was zero stations per night during the 2020 summer. Night-time ozone has been considered when hourly concentration has exceeded the level $100 \mu\text{g}/\text{m}^3$ between 10PM and 2AM at any station. Night-time ozone is most frequently in Gavalinagar, Pimpri and Savitribai Phule University, Pune where it was reported on 28 nights and 26 nights respectively (See *Table 5: Locations with most night-time ground-level ozone pollution instances*).

Ground-level ozone has become a yearlong problem: Even though the ground-level ozone exceedance is the worst during summer months, it remains a year-long problem as at least few locations continue to record exceedance throughout the year. The dangerous build-up of ground-level ozone can happen anytime during the year, but it is usually in small pockets during non-summer months. For it to have wider spatial spread hot and sunny weather conditions are needed which are generally present in Pune throughout the year outside monsoon. Ground ozone pollution in Pune is at its worse during April-May. Exceedances go down from May onwards and then there is a considerable uptick in ozone pollution after monsoon – especially October-November (See *Graph 5: Trend in ground-level ozone over years -2020-24*). Considerable number of exceedance are recorded during January-March period as well. This year so far 138 days of exceedance have been recorded in Pune Metropolitan Region.

Graph 4: Daily variation in occurrence of high night-time ground-level ozone pollution



Note: Based on high hourly concentration of ground-level ozone recorded at the monitoring stations in Pune Metropolitan Region during night-time. High hourly concentration is taken as 100 µg/m³ or more. Night-time is taken as 10PM to 2AM. Period of study is 1 April to 18 July 2024.

Source: CSE analysis of CPCB realtime data.


Table 5: Locations with most night-time ground-level ozone pollution instances

SNo	Station	Number of night-time exceedance
1	Gavalinagar, Pimpri	28
2	Savitribai Phule University, Pune	26
3	Karve Road, Pune	23
4	Thergaon, Pimpri	20
5	Alandi, Pune	11
6	Mhada Colony, Pune	10
7	Bhosari, Pune	5
8	Katraj Dairy, Pune	5
9	Shivajinagar, Pune	4
10	Ngdi, Pune	3
11	Park Street Wakkad, Pimpri	3
12	Hadapsar, Pune	0
13	MT, Pune	0
14	Bhunkar Nagar, Pune	0
15	Panchawati, Pune	0

Note: Based on high hourly concentration of ground-level ozone recorded during night-time. High hourly concentration is taken as 100 µg/m³ or more. Night-time is taken as 10PM to 2AM. Period of study is 1 April to 18 July 2024.

Source: CSE analysis of CPCB realtime data.

Graph 5: Trend in ground-level ozone over years -2020-24



Note: Based on exceedances recorded at the monitoring stations in Pune Metropolitan Region. Exceedance is computed as daily maximum 8-hr average crossing the ground-level ozone 8-hr standard, i.e. 100 µg/m³. Duration of exceedance is computed as number of hours the rolling 8-hr average was exceeded at a station on a day. Period of the study is 1 April to 18 July.
Source: CSE analysis of CPCB realtime data.

Act now

Ozone mitigation demands stringent control of gases from all combustion sources including vehicles, industry, power plants and open burning in the entire region. It is therefore necessary that while designing mitigation of particulate matter the key focus of action strategy today, is also calibrated for reduction of ozone precursor gases.

Immediately, refine the action strategy for combined control of particulate pollution, ozone and its precursor gases like NO_x to maximise the co-benefits of the action plan.

Simultaneously develop a robust public information and dissemination system to alert public about ozone exceedance wherever ozone build up is happening for exposure management.